


**Marine litters monitoring system using Earth
Observation, COPERNICUS models and
citizen/participative science**

Antoine Mangin^{1,2}, Francois-Regis Martin-Lauzer¹,
Manuel Arias¹ Mark Hennen¹, James Delaney¹

Outlines of this presentation

- Marine litters – the situation (focus in the Med)
- Needs for observation
- Means for observation (ARGANS perspective)
 - Earth Observation
 - Modelling
 - [Citizen/participative science](#)
- Conclusions - Next steps





Out of sight, out of mind






Figure 4.--Densities of fragments, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 288,000 pieces/km².

- ✓ No evidence of garbage patch in the Med – today
- ✓ Mainly plastics



- ✓ Estimates : 62 millions of macrowaste floating for the whole Med
- ✓ Everything is coming from land and sea (even through Gibraltar) – no escape.

Biological and Eco-societal impacts



Environmental regulations

- Marine Strategy Framework Directive (marine litter is descriptor #10)
- Mediterranean Action Plan (Barcelona convention – UNEP) – Regional Plan 2014
- ...

Needs to monitor locations of:

1. Emission / release of macro-waste (for reduction/ control)
2. Landing/beaching (for pollution fight, coastal spatial zoning, moorings and navigation routes)
3. Floating filaments and patches (for cleaning)

Monitoring

Emission / release of macro-waste

- EO (sentinel-2)
- Modelling (emission coefficient based on land use on watershed)
- Participative observation by professional

Landing/beaching

- EO (sentinel-2)
- Modelling (statistics)
- Participative observation by citizens and professional

Floating filaments and patches

- EO (sentinel-2 / Sentinel-1 ?)
- Participative observation by marine professions (fishermen, transportation...)

Objective is to merge efforts to demonstrate what is feasible/useful and could be operationally deployed -> **consolidate a roadmap in addition to existing efforts.**



Marine Litter Programme



Remote Sensing for Marine Litter
-RESMALI-



GeoInt Service for Marine Litter
-SSGP Project-





EO Tracking of Marine Litter in
the Mediterranean Sea



Copernicus Service for
Marine Litter

Goals of the RESMALI project

- Characterization of physical properties and meaningful spatial and time scales for marine litter.
- Definition of remote sensing observational requirements of data acquisition for scientific and downstream applications.
- Identification of best potential technologies and instrument ensemble that could compose an EO mission for marine litter.



Characterization of marine litter

➤ Based on the **marine domain** under consideration



Beaches



Shallow waters



Open Ocean

➤ Based on the **marine litter fraction** under observation



Large Items



Small pieces and fragments



Microplastics

Characterization of marine litter

ML Characteristic	Domain			
	Open Ocean	Continental Shelf	Coastal Area	Shores & Beaches
Variability	Low (months to years)	Medium (months to weeks)	Medium to High (weeks to days)	Very high (days to hours)
Residence time	Long	Medium to short	Short	Long to short
Origin	Plastics (PP, PE)	Plastics (PP, PE, PS, PET) Organic	Plastics (PP, PE, PS, PET) Organic, Rubber, Paper	Plastics (PP, PE, PS, PET) Organic, Rubber, Paper, Metal
Accumulation factor	Global currents	Wind/currents transport, human activity	River mouths, run-offs, dumping, human activity	Tidal/storm stranding, dumping, human activity

Complexity



Characterization of marine litter

Characteristic	ML fraction			
	> 200mm	200-5mm	5-1mm	<1mm
Abundance	Very low	Low to medium	Very high	?
Total mass	Very high	High to medium	Medium to very low	?
Vertical zoning	0-5m	0-5m	0-5m	?
Main composition	PE, PP, PET, PS	PE, PP, PET, PS	PE, PP	PE, PP

Large items accumulate most of mass but are the less abundant

Smallest fractions are composed by fewer materials

Most of observable ML is found at the surface of the oceans


Deployment of 10x10m patches of plastic materials in Lesbos Island. Source: K. Topouzelis



Project: EO Tracking of Marine Debris in the Mediterranean Sea

Deployment of 10x10m patches of plastic materials in Lesbos Island

Sentinel-2/MSI Image Natural colour RGB (Band 4 / 665 nm, Band 3 / 560 nm, Band 2 / 490 nm)




02/02/2018

Sentinel-2/MSI has capability to observe sub-pixel marine litter


Deployment of 10x10m patches of plastic materials in Lesbos Island

Sentinel-2/MSI Image Natural colour RGB (Band 4 / 665 nm, Band 3 / 560 nm, Band 2 / 490 nm)



Selection of areas for spectral data study

Deployment of 10x10m patches of plastic materials in Lesbos Island Sentinel-2/MSI spectral data over selected classes



Plastic litter has identifiable pikes in NIR (Band 8 – 835 nm) and SWIR (Band 11 – 1613 nm)




Mithi River, Mumbai

Plastic pollution along the raised banks of the Mithi River, Mumbai. Top left: Google Earth image showing the modified section of Mithi River (Top right). Source: Hindustan Times.




23/04/2018



19/03/2018


Mithi River garbage patches (red polygon) as seen in Sentinel-2/MSI Natural color RGB image
Sentinel-2 can capture variability of litter in targeted and confined areas




S-2/MSI Spectral profiles for Mithi River

However, low concentrations of litter are much more difficult to detect, whereas scattered in water (purple line) or mixed with vegetation (red line)


Dispersion of a generalised, uniform and unitary pollution of floating objects
After 12 hours of drift




After 24 hours




After 36 hours




After 48 hours




After 60 hours



Beaching/Landing





Simplex **ACRI**
HE

Automated tool that generates colocated and coregistered information to improve knowledge communication and science

Here an elegant and easy mixing of :

- Participative observations
- Modeling at large scale (me...)
- EO at regional scale (SST...)


This has been adapted to marine litters With an updated application and webGIS


La... (2018) : **6400 observers**
Today the jellyfish database has **34000 records**



Legend items: e-Delphis, Dordogne, Foret, Seaplex, Relax, Méduses, Données environnementales, **Température de surface de la mer**, SECCHI, CHLA, Air, Meteo, Nuage, Pluie, Vent, Humidité relative, Température de surface, Terre.

Map layers: Vent (0-30ms), SST (0-28°C).

Timeline: Août, Sept.







...and the map

Monitoring with « citizen » - is not only a matter of tool !

The most difficult part is to mobilise people – under which conditions and to which level of participants ?

Emission / release of macro-waste

- Participative observation by professional network of river monitoring and citizens (environmental association) for Wadi (traps of macrowaste)

Landing/beaching

- Participative observation by citizens and professional (many associations are existing, contacts are taken in France, Morocco , Algeria) – organisation of observation in // with « the annual cleaning day »

Floating filaments and patches

- Participative observation by marine professions (fishermen, transportation...)

The monitoring system for marine litters in the Western Mediterranean is built around different and complementary facets of the observation (the *NewMonitoring* ?)

Use of EO

Interesting results for specific spectral bands (or pairs of) for marine litter detection
Involvement of new actors in the field that would also help the participative observations (e.g. the ocean clean up)

Citizen/ participative observatories

Reliable tool for monitoring

Engagement of the participants :

- Make the distinction between : one-shot campaign every year and continuous observation,
- Rewarding participant ?
- Making them participate to the final analysis...

Modelling is beginning

Thank you for attention


**Marine litters monitoring system using Earth
Observation, COPERNICUS models and
citizen/participative science**

**Antoine Mangin^{1,2}, Francois-Regis Martin-Lauzer¹,
Manuel Arias¹ Mark Hennen¹, James Delaney¹**



Polygons

Control Bar (layers)




Time Silder

Curent location

Zoom In/Out

Exemple of observation of today

GéoSimplex Simplex plus + Download + ARGANS




The screenshot shows the GeoSimplex web application interface. The main map displays a portion of Italy, with a popup window centered over a location. The popup contains a photograph of an interior space, likely a meeting room or office, and a table with the following data:

Id Obs	28
Observation	Clothes/textile
Lon	12.6741
Lat	41.8264
Alt (m)	231.000
Date/Heure (GMT+0)	14/10/2018 14:24:28

On the right side of the map, there is a legend titled "Observé" with the following categories and colors:

- Processed wood (Yellow)
- Paper/Cardboard (Blue)
- Plastics (Pink)
- Don't know (Magenta)
- Metal (Red)
- Glass/Ceramics (Yellow)
- Clothes/textile (Black)

At the top left, there are navigation icons for zooming and a scale bar. At the top right, there are map style options: "Street Map" (selected) and "Aerial Imagery".



- Simplex Gallerie
- Simplex Stat


Simplex Gallery

Collecte test


13/11/2018 12:11:11

13/11/2018 12:11:36

13/11/2018 12:11:57



13/11/2018 12:11:38



13/11/2018 12:11:56



14/11/2018 12:11:51



14/11/2018 14:11:28




LA River garbage boom

Source: Google Earth. (Top right) View from the western river bank looking east over the garbage boom during dredging (Unknown date / source). (Bottom right) View from the eastern river bank over the lagoon (Golden Shore Marine Biological Reserve). Source: Google Earth Street view.




LA River garbage boom Sentinel-2/MSI Natural color RGB image (02/02/2018)

Litter patch observed in the expected location (red polygon)




LA River garbage boom Sentinel-2/MSI Natural color RGB image (05/03/2018)

Litter patch observed in the expected location (red polygon)



02/02/2018



05/03/2018

S-2/MSI Spectral profiles for L. A. River

Litter (red line) has a differential spectra with respect water (blue) and vegetation (green)